

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :

A61K 9/00, 47/18

A1

(11) International Publication Number:

WO 99/24015

(43) International Publication Date:

20 May 1999 (20.05.99)

(21) International Application Number: PCT/US98/23411

(22) International Filing Date: 3 November 1998 (03.11.98)

(30) Priority Data:

08/969,217

12 November 1997 (12.11.97) US

(71) Applicant: ALZA CORPORATION [US/US]; 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US).


(72) Inventors: LEUNG, Iris, Ka, Man; 14727 Greenleaf Valley Drive, Chesterfield, MO 63017-5514 (US). CORMIER, Michel, J., N.; 278 Andsbury Avenue, Mountain View, CA 94043 (US). SENDELBECK, Sara, Lee; 3505 Laguna Avenue, Palo Alto, CA 94306-2631 (US). MUCHNIK, Anna; 1542 Winding Way, Belmont, CA 94002 (US).

(74) Agents: MILLER, D., Byron et al.; Alza Corporation, 950 Page Mill Road, P.O. Box 10950, Palo Alto, CA 94303-0802 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published*With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: BUFFERED DRUG FORMULATIONS FOR TRANSDERMAL ELECTROTRANSPORT DELIVERY



(57) Abstract

Buffered drug formulations for transdermal electrotransport delivery are disclosed. The formulations utilize a dipeptide as a buffer and allow for more efficient electrotransport delivery of drugs, e.g., polypeptide drugs, via the transdermal route.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

1 Claims:

2 1. A formulation for transdermal electrotransport delivery,
3 comprising an aqueous solution of a drug or an electrolyte and a dipeptide
4 buffer, the dipeptide buffer comprising a polypeptide chain of 2 to 5 amino
5 acids and having an isoelectric pH at which the dipeptide carries no net
6 charge, the dipeptide having at least 2 pKa's which are separated by no more
7 than about 3.5 pH units, the solution having a pH which is within about 1.0 pH
8 unit of the isoelectric pH.

9
10 2. The formulation of claim 1, wherein the isoelectric pH of the
11 dipeptide is between about 3 and 10.

12
13 3. The formulation of claim 1, wherein the dipeptide is present
14 in the solution at a concentration of at least about 10 mM.

15
16 4. The formulation of claim 1, wherein the dipeptide includes at
17 least one amino acid selected from the group consisting of His, Tyr, Arg, Cys,
18 Lys, Asp and Glu.

19
20 5. The formulation of claim 1, wherein the dipeptide includes
21 His.

22
23 6. The formulation of claim 1, wherein the dipeptide is Gly-His.

24
25 7. The formulation of claim 1, wherein dipeptide is selected
26 from the group consisting of Asp-Asp, Gly-Asp, Asp-His, Glu-His, His-Glu,
27 His-Asp, Glu-Arg, Glu-Lys, Arg-Glu,, Lys-Glu, Arg-Asp, Lys-Asp, His-Gly, His-
28 Ala, His-Asn, His-Citruline, His-Gln, His-Hydroxyproline, His-Isoleucine, His-
29 Leu, His-Met, His-Phe, His-Pro, His-Ser, His-Thr, His-Trp, His-Tyr, His-Val,
30 Asn-His, Thr-His, Try-His, Gin-His, Phe-His, Ser-His, Citruline-His, Trp-His,
31 Met-His, Val-His, His-His, Isoleucine-His, Hydroxyproline-His, Leu-His, Ala-

1 His, Gly-His, Beta-Alanylhistidine, Pro-His, Carnosine, Anserine, Tyr-Arg,
2 Hydroxylysine-His, His-Hydroxytyrosine, Ornithine-His, His-Lys, His-Ornithine
3 and Lys-His.

4
5 8. The formulation of claim 1, wherein the drug comprises a
6 polypeptide or a protein.

7
8 9. A transdermal electrotransport drug delivery device (10)
9 having a reservoir (26, 28) containing the formulation of claim 1.

10
11 10. A transdermal electrotransport drug delivery device (10)
12 having a drug-containing donor reservoir (26, 28) containing the formulation
13 of claim 1.

14
15 11. A transdermal electrotransport drug delivery device (10)
16 having a electrolyte-containing counter reservoir (26, 28) containing the
17 formulation of claim 1.

18
19 12. A method of buffering an aqueous solution of a drug or an
20 electrolyte used for transdermal electrotransport delivery, comprising buffering
21 the solution with a dipeptide comprising a chain of 2 to 5 amino acids and
22 having an isoelectric pH at which the dipeptide carries no net charge, the
23 dipeptide having at least 2 pKa's which are separated by no more than about
24 3.5 pH units, the solution having a pH which is within about 1.0 pH unit of the
25 isoelectric pH.

26
27 13. The method of claim 12, wherein the isoelectric pH of the
28 dipeptide is between about 3 and 10.

29
30 14. The method of claim 12, wherein the dipeptide is present in
31 the solution at a concentration of at least about 10 mM.

1 15. The method of claim 12, wherein the dipeptide contains one
2 or more of His, Tyr, Arg, Cys, Lys, Asp and Glu.

3
4 16. The method of claim 12, wherein the dipeptide contains His.

5
6 17. The method of claim 12, wherein the dipeptide is Gly-His.

7
8 18. The method of claim 12, wherein dipeptide is selected from
9 the group consisting of Asp-Asp, Gly-Asp, Asp-His, Glu-His, His-Glu, His-Asp,
10 Glu-Arg, Glu-Lys, Arg-Glu,, Lys-Glu, Arg-Asp, Lys-Asp, His-Gly, His-Ala, His-
11 Asn, His-Citruline, His-Gln, His-Hydroxyproline, His-Isoleucine, His-Leu, His-
12 Met, His-Phe, His-Pro, His-Ser, His-Thr, His-Trp, His-Tyr, His-Val, Asn-His,
13 Thr-His, Try-His, Gin-His, Phe-His, Ser-His, Citruline-His, Trp-His, Met-His,
14 Val-His, His-His, Isoleucine-His, Hydroxyproline-His, Leu-His, Ala-His, Gly-
15 His, Beta-Alanylhistidine, Pro-His, Carnosine, Anserine, Tyr-Arg,
16 Hydroxylysine-His, His-Hydroxytlysine, Ornithine-His, His-Lys, His-Ornithine
17 and Lys-His.

18
19 19. The method of claim 12, wherein the drug comprises a
20 polypeptide or a protein.

21
22 20. The method of claim 12, wherein the solution is contained
23 in a reservoir of a transdermal electrotransport drug delivery device.

24

25

26

1/11




FIG. 1

2/11




FIG. 2

3/11




FIG. 3

4/11




FIG. 4

5/11




FIG. 5

6/11




FIG. 6

7/11




FIG. 7

8/11

FIG. 8






FIG. 9





11/11




FIG. 11